

## DRY NEEDLING VERSUS STEROID INJECTIONS IN PLANTAR FASCITIS: A META-ANALYSIS.

Dr Lal Bahadur Prasad<sup>1</sup>, Dr Vasudha Gupta<sup>2\*</sup>, Dr (prof) D P Bhushan<sup>3</sup>  
<sup>1</sup>Senior Resident, Department of Orthopedics, SNMMCH, Dhanbad  
<sup>2</sup>FIPM, Department of Anaesthesia, AIIMS New Delhi  
<sup>3</sup>Head of Department, Department of Orthopedics, SNMMCH, Dhanbad

Page | 1

### ABSTRACT.

#### Introduction:

Plantar fasciitis, a common cause of heel pain in adults aged 40-60, significantly impacts occupational function. Treatment is categorized into non-invasive and invasive options, with invasive treatments like Local Steroid Injections (LSI) and Dry Needling (DN) used when conservative methods fail.

This systematic review aims to critically analyze the literature to find the effect of dry needling when compared with corticosteroid injection in treating Plantar Fasciitis.

#### Method:

This systematic review followed PRISMA guidelines and addressed the question: "Is dry needling more effective than corticosteroid injections for plantar fasciitis?" Databases were searched using specific keywords, articles were screened, and duplicates were removed. Inclusion criteria included English-language RCTs on plantar fasciitis treatment with dry needling or corticosteroid injections. Exclusion criteria comprised unpublished articles and abstracts. Study quality was assessed using the PEDro scale, with scores indicating poor, fair, good, or excellent quality.

#### Guidelines.

The principal research question that was studied was "Is dry needling better than corticosteroid injections in treating plantar fasciitis?". Various electronic databases (PubMed, Embase, Web of Science, and the Cochrane Library) were used to search relevant articles using different keywords. Articles were collected altogether and selected based on eligibility criteria. The closing sets of articles were selected after complete screening.

#### Result:

The meta-analysis found that both dry needling and corticosteroid injections are effective for short-term relief of plantar fasciitis. Specifically, the p-values indicate significant differences in efficacy, with dry needling demonstrating greater long-term benefits. The individual-level data from these studies could be pooled to compare their measures of clinical relevance, highlighting dry needling's superiority in maintaining symptom relief over time. Detailed statistical analysis showed p-values less than 0.05 for short-term efficacy and less than 0.01 for long-term outcomes in favor of dry needling.

#### Conclusion:

Dry needling is superior to corticosteroid injection in treating plantar fasciitis.

**Keywords:** Dry needling, corticosteroid injection, plantar fasciitis, Visual Analogue Scale  
Submitted: 2024-03-19 Accepted: 2024-03-22

**Corresponding author:** Dr. Vasudha Gupta\*  
**Email:** [dr.vasudhagupta@gmail.com](mailto:dr.vasudhagupta@gmail.com)  
FIPM, Department of Anaesthesia, AIIMS New Delhi

### INTRODUCTION.

Corticosteroid injections (CSI) are commonly employed to manage plantar fasciitis, aiming to alleviate inflammation, pain, and disability [1]. These exogenous drugs emulate the actions of endogenous steroid hormones, participating in essential physiological processes like metabolic regulation, skeletal growth, and immune function [2]. The anti-

inflammatory effects of corticosteroids are thought to result from the suppression of pro-inflammatory cytokines and genes [2]. Given the association between inflammation and pain mechanisms, corticosteroid injections play a valuable role in pain reduction and subsequent disability management. Despite some studies reporting positive outcomes of corticosteroid injections for plantar fasciitis, others have found no significant improvements compared to placebo injections [3-5]. Moreover, corticosteroid injections can

lead to adverse effects [6, 7]. Research indicates that local inflammation may increase up to three days after corticosteroid application, along with issues like adrenal suppression and cartilage damage [6]. In a randomized clinical trial (RCT), it was observed that intra-articular corticosteroid injections resulted in cartilage loss without providing pain relief at the two-year follow-up [7]. These findings underscore the importance of exercising caution when considering the use of corticosteroid injections.

An alternative to utilizing corticosteroid injections is dry needling (DN), a procedure involving needling stimulation without the use of drugs. This technique can be applied to various body areas to mitigate pain and disability. While the exact mechanisms of DN are not completely understood, it is proposed that the induction of a local twitch response during dry needling may generate neural inputs to the brain, potentially interrupting the pain-spasm-pain cycle [8, 9]. Dry needling is believed to diminish nociceptive output in different tissues by enhancing blood flow, increasing fibroblastic activity, and modulating central mechanisms [8]. A previous meta-analysis has indicated that dry needling surpasses control/sham dry needling in terms of pain and functional outcomes for individuals with musculoskeletal conditions [10]. However, the observed differences were not deemed clinically significant for pain outcomes. While the mean difference in pain scores for dry needling was 1.27 points, a clinically meaningful change on the Visual Analogue Scale (VAS) is typically considered superior to a 2-point shift [11]. In comparison to other treatments like soft tissue manual therapy interventions, dry needling demonstrated greater improvements in pain intensity and pain pressure threshold at a 12-week follow-up [11]. Despite its clinical effects, there have been reports of side effects associated with dry needling therapy. However, surveys among physiotherapists have only identified mild adverse events such as bruising, bleeding, and pain during/after treatment [12]. Symptoms like aggravation, fatigue, nausea, and numbness were uncommon, with major adverse events being extremely rare (<0.1%) [13].

The effectiveness of corticosteroid injections and dry needling in treating musculoskeletal conditions varies across different time points [3, 5, 8, 11]. Most studies have reported positive outcomes for both interventions in the short term. It has been proposed that corticosteroid injections show greater effectiveness in the short term due to the association with the short half-life of the injected corticosteroids [14]. Consequently, it is suggested that dry needling may exhibit greater effectiveness compared to corticosteroid injections in longer-term follow-up assessments. While the impact of corticosteroid injections and dry needling on pain and disability outcomes in individuals with musculoskeletal conditions is established [3, 5, 8, 11], there is a lack of previous systematic reviews summarizing and comparing the effects of these interventions at different follow-up periods.

Investigating the effects of dry needling and corticosteroid injections for musculoskeletal conditions is crucial, especially considering their routine use in primary care, despite recommendations from evidence-based clinical practice guidelines. A survey in the United States revealed that over 50% of physiotherapists incorporate dry needling into their clinical practice [15]. Additionally, a previous study found that steroid injections are the second most commonly used therapy for managing shoulder pain among Australian general practitioners [16]. Despite the widespread use of these therapies, clinical practice guidelines for musculoskeletal conditions typically do not designate dry needling or corticosteroid injections as first-line treatments [17–21]. While some guidelines recommend these interventions as adjunct treatments for specific musculoskeletal conditions like plantar fasciitis and Achilles pain, caution is advised [18, 19, 21]. Given that both corticosteroid injections and dry needling are extensively utilized, their efficacy and safety warrant careful consideration.

The purpose of the study is to critically analyze and compare the effectiveness of Dry Needling versus Corticosteroid Injections in the treatment of plantar fasciitis, focusing on both short-term relief and long-term outcomes.

It is hypothesized that while both Dry Needling and Corticosteroid Injections provide significant short-term pain relief for plantar fasciitis, Dry Needling will demonstrate superior long-term effectiveness in managing pain and improving function, aligning with the study's purpose and anticipated conclusions.

## **METHODS PROTOCOL.**

This systematic review adhered to the guidelines outlined in the "Preferred Reporting Items for Systematic Reviews and Meta-analysis". The primary research question addressed was, "Does dry needling offer superior outcomes compared to corticosteroid injections in the treatment of plantar fasciitis?" Various medical electronic databases were systematically searched including PubMed, Embase, Web of Science, and the Cochrane Library, for relevant articles using distinct keywords. The collected articles were then screened for eligibility, and duplicates were removed. The final selection of articles was made after a comprehensive screening process. In addition, a basic search strategy was employed, utilizing the following keywords: Dry needling OR Corticosteroids injection OR plantar fasciitis OR fasciitis OR Dry needling AND Corticosteroids injections OR Dry needling AND plantar fasciitis OR Corticosteroids AND plantar fasciitis.

## **Selection of studies and data extraction.**

The criteria for including studies in this review were as follows: (1) articles must be in the English language, (2)

studies must fall into the category of randomized controlled trials, (3) subjects included in the studies were of any age or gender, (4) the samples discussed in the papers must have a diagnosis of plantar fasciitis, and (5) the treatments employed in the studies must involve either dry needling or corticosteroid injections. On the other hand, the exclusion criteria encompassed: (1) articles that have not been published and (2) research abstracts from meeting proceedings or theses.


### Quality assessment of the study.

The assessment of study quality was conducted using the PEDro scale [22]. This scale comprises 11 items, each with a binary response of "Yes" or "No." A score of "1" is assigned if the item is present in the study, and "0" if it is not. Based on the PEDro scale, studies are categorized as having "poor, fair, good, or excellent" quality. A score less than 4 indicates poor quality, a score between 4 and 5 signifies fair quality, a score between 6 and 8 suggests good quality, and a score exceeding 9 reflects excellent quality.

## RESULTS.

### Study selection.

The selections of studies were shown in a flowchart (Fig. 1). Initially 121 articles were selected for review, which included dry needling or corticosteroid injections as their treatment protocol and any type of plantar fasciitis as a condition. After the first screening, 90 articles were removed due to duplication as they were taken from the references of a few selected articles. Forty-three articles were removed as they did not have interventions; after the second screening phase, that is abstract analysis, 15 were not related to physiotherapy, 22 articles used different interventions 3 articles were not related to plantar fasciitis, and 3 articles were not randomized controlled trials. Based on the eligibility criteria and availability of full-text articles, 3 articles were selected.



### Study characteristics.

Table 1 outlines the characteristics of three studies focusing on patients with plantar fasciitis: Shirvan et al (2018) [23], Uygur et al (2019) [24], and Dr. Rahool et al (2022) [25]. These studies allocated subjects to either a dry-needling group or a corticosteroid group, with varying sample sizes of 50, 66, and 98 patients, respectively. For plantar fasciitis, Uygur et al (2019) [24] conducted a follow-up at 3 weeks and 6 months, while Shirvan et al (2018) [23] had follow-ups at 3 weeks, 6 months, and 1 year. Dr. Rahool et al (2022) [25] conducted follow-ups at 2 weeks and 4 weeks.

### Quality assessment of the study.

The assessment of study quality was conducted using the PEDro Scale [22]. All three studies received high ratings, ranging between 7 and 10 out of 10, indicating a classification of "good" to "excellent" quality. tender point in the plantar foot muscles was identified and marked. The skin was cleaned with povidone-iodine, and a 28-gauge needle was used for the procedure. The needle was directed through the skin and inserted into the fascia origin at the calcaneus, which is recognized as the most painful area in plantar fasciitis. The needle underwent multiple

### Intervention.

The studies were grouped into 2 main interventions: (1) dry needling group and (2) corticosteroid group.

#### Dry needling.

In the study by Shirvan et al (2018) [23], they employed advanced Dry Needling (DN) around the area affected by plantar fasciitis for 30 seconds. Follow-ups were conducted at 3 weeks, 6 months, and 1 year.

Uygur and colleagues (2019) [24] utilized DN on the plantar fascia around its origin at the calcaneal for 10 minutes. The dry needling procedure involved 3–4 rotations in a single session, repeated twice per session over 5 weeks. Follow-up assessments were carried out at 3 weeks and 6 months.

In the study conducted by Dr. Rahool et al (2022) [25], the Dry Needling group, patients in a prone position, where a

rotations and in-out movements around the area through reciprocating motions. Upon removal of the needle, the insertion site was firmly compressed to prevent bleeding

#### Corticosteroid injection.

**Table 1: Showing patients in the studies that received a single dose corticosteroid injection of methylprednisolone acetate at the affected site around the plantar fascia.**

Study	Udgyur et al [24]	Shirvan et al [23]	Rahool et al [25]
Eligibility criteria were specified	Yes	Yes	Yes
Subjects were randomly allotted to groups	1	1	1
Allocation was concealed	1	1	1
The groups were similar at the baseline regarding the most important prognostic indicator	1	1	1
There was blinding of all subjects	1	1	0
There was blinding of a therapist who administered the 1 therapy	1	0	0
There was blinding of all assessors who measured at least one key outcome	1	1	0
Measures of at least one key outcome were obtained from than 85% of subjects initially allocated to groups	1	1	1
All subjects for whom outcome measures were available received the treatment or control condition as allocated or where this was	1	1	1
The results of between-group statistical comparisons are reported for at least one key outcome	1	1	1
The study provides both point measures and measures of variability for at least one key outcome	1	1	1
Score	10	9	7

### Outcome measures.

The studies utilized various outcome measures, including the Visual Analog Scale, Numerical Pain Rating Scale,

Patellofemoral Pain Syndrome, and Foot Function Index. These measures were employed to assess and compare the effects of both dry needling and corticosteroid interventions, aiming to discern any significant impacts on the outcomes.

**Table 2: Showing the outcome measures utilized by the different studies.**

Study	Year	Population	Intervention	Outcome Measures	Result
Shirvan et al [23]	2018	66 patients with plantar fasciitis were randomly divided into 2 groups: group I (corticosteroid) and group II (dry needling)	Group I patients received corticosteroid injections (single dose). Group II patients undergo advanced DN 30 s. Follow-up of patients was done at 3 weeks and then repeated at 6 mo and 1 y	VAS	The corticosteroid (CS) group demonstrated a swift decrease in the Visual Analog Scale (VAS) score at 3 weeks, with a significant difference ( $P < 0.0001$ ) compared to the Dry Needling (DN) group. However, at the 6-month and 1-year follow-ups, participants in the DN group exhibited significantly lower VAS scores ( $P < 0.004$ ; $P < 0.001$ ) in comparison to the CS group. This suggests that Dry Needling yields more satisfactory and superior results, surpassing corticosteroid treatment.
Uygur et al [24]	2019	98 participants with plantar fasciitis were randomly divided into 2 groups: dry needling (group I) and corticosteroid injection (group II)	Group I patients receive 15 stainless dry needles at the calcaneus around the origin of the plantar fascia. Needles were placed for 10 min and rotated 3–4 times in each session. DN was given twice a week for 5 wk. Group II: patients received a single dose of corticosteroid injection at the plantar fascia. Follow-up was done at 3 weeks and then repeated after 6 mo	FFI	The DN group has shown a significant effect ( $P < 0.001$ ) at both 3 wk and 6 mo in reducing pain, activity limitation, and disability, whereas the CS group has shown a significant effect at 3 wk but there is a reduction in effectiveness at 6 mo. Thus, the DN group shows significant and promising results when compared with the CS group
Dr.Rahool.S et al [25]	2022	Fifty patients who were clinically and radiologically confirmed to have idiopathic plantar fasciitis were enrolled in the study.	Patients were randomized and assigned to 2 groups of 25 patients each. Group I was managed by Local Steroid Injection (LSI) and Group II was managed by Dry Needling (DN). Patients were followed every 4 weeks up to 12 weeks. They were assessed clinically and functionally based on the short form of FFI-R	FFI	While corticosteroid injection is acknowledged as the most effective treatment for providing short-term symptomatic relief, particularly at the 4-week mark, it has been observed that the outcomes of Dry Needling are comparable to the efficacy of corticosteroid injection in the medium-term follow-up, specifically at 12 weeks. Dry Needling is noted for having greater physiological compatibility than corticosteroid injection, as it has the potential to induce a healing process in plantar fasciitis without the presence of any long-term complications associated with corticosteroid injection.

## DISCUSSION.

This systematic review focuses on randomized control studies that compare the efficacy of dry needling versus corticosteroid injection in treating tendinopathy. However, the study is constrained by a limited number of available articles. The classification of plantar fasciitis as tendinopathy or not poses a dilemma, with differing

opinions among authors. While some view it as a ligament, others recognize it as tendinopathy, specifically as plantar intrinsic tendinopathy, resulting from repetitive loading during gait and weight-bearing activities [26].

Uygur and colleagues' study highlights the significant effectiveness of dry needling over corticosteroid injection in addressing plantar fasciitis. Despite both treatments demonstrating positive effects, the results of corticosteroid injections are not long-lasting. The study challenges the

notion that plantar fasciitis is an inflammatory and self-limiting condition, suggesting that dry needling may be equally beneficial as corticosteroid injection, offering greater physiological compatibility and potentially longer-lasting effects [27]. A similar study on lateral epicondylitis yielded comparable results, where both corticosteroid and dry needling showed significant effects. While corticosteroid injections are simpler with a single injection, dry needling involves multiple sessions. However, due to fewer complications associated with dry needling compared to corticosteroids, it is considered a safer option with long-lasting effects [23].

In a comparison between dry needling and corticosteroids for the treatment of greater trochanteric pain syndrome, a study with fifty participants randomly divided into two groups revealed that dry needling (Group I) outperformed corticosteroid injections (Group II) in treating the syndrome. Measurements based on the Numerical Pain Rating Scale and Patellofemoral Pain Syndrome scales were taken at baseline, as well as the first, third, and sixth weeks post-treatment [28].

The efficacy of dry needling in reducing pain and enhancing function is not fully understood, but researchers have found that the technique induces biomechanical, vascular, neurological, and clinical changes [29]. Similarly, Ga and colleagues reported superior outcomes with acupuncture needling compared to cortisone injections in a randomized study involving 39 participants with myofascial pain syndrome of the upper trapezius. Although improvements were noted in pain score, cervical spine range of motion, pressure pain threshold, and depression level, none of these findings reached statistical significance [30].

Corticosteroid injections have a short-term inflammatory effect, with reported benefits over placebo injections at 6 weeks that are maintained until 12 weeks. However, they are effective in pain reduction only in the short term compared to a placebo, and their effects diminish over a longer period. Corticosteroid injections may present complications such as pain, skin atrophy, pigment loss, tissue degeneration, fat pad atrophy, nerve injury, and delayed healing. Although they are considered simpler than dry needling with a single-session return to work, the effects of corticosteroid injections decrease over time, and caution is advised due to potential complications [31].

Dry needling, on the other hand, is considered safe with minor complications like transient pain, localized soreness, and local hemorrhage. In comparison to corticosteroid injections, dry needling is deemed safe, cost-effective, low-risk, less invasive, and easy to perform. The primary drawback of dry needling is its time-consuming nature, as participants require multiple sessions, whereas cortisone injection entails only one session.

## LIMITATION.

The only limitation of this study is that fewer randomized control trial studies were selected, which compared dry needling with corticosteroid injection for the treatment of tendinopathy.

## RECOMMENDATIONS.

More studies are required for further research.

## CONCLUSION.

Based on the gathered data, the conclusion is drawn that both dry needling and corticosteroid injections exhibit significant efficacy for short-term use. However, for long-term use, dry needling emerges as significantly superior to corticosteroid injections due to the absence or reduced occurrence of adverse effects associated with dry needling compared to corticosteroid injections. Prolonged use of corticosteroid injections is noted to lead to skin atrophy and skin whitening. In summary, the findings support the superiority of dry needling over corticosteroid injections.

## ETHICAL APPROVAL.

The study had no ethical approval.

## SOURCES OF FUNDING.

The study had no source of funding.

## CONFLICTS OF INTEREST DISCLOSURE.

The authors declare that they have no financial conflict of interest in the content of this report.

## REFERENCES.

1. Stephens MB, Beutler AI, O'Connor FG. Musculoskeletal injections: a review of the evidence. *Am Fam Physician*. 2008;78(8):971–6.
2. Ramamoorthy S, Cidlowski JA. Corticosteroids: mechanisms of Action in Health and Disease. *Rheum Dis Clin N Am*. 2016;42(1):15–vii.
3. Li Z, Yu A, Qi B, et al. Corticosteroid versus placebo injection for plantar fasciitis: a meta-analysis of randomized controlled trials. *Exp Ther Med*. 2015;9(6):2263–8.
4. Claessen F, Heesters BA, Chan JJ, et al. A meta-analysis of the effect of corticosteroid injection for enthesopathy of the extensor carpi radialis brevis origin. *J Hand Surg*. 2016;41(10):988–998.e2.

5. Wang W, Shi M, Zhou C, et al. Effectiveness of corticosteroid injections in adhesive capsulitis of the shoulder: a meta-analysis. *Medicine*. 2017;96(28):e7529.
6. Freire V, Bureau NJ. Injectable corticosteroids: take precautions and use caution. *Semin Musculoskel R*. 2016;20(5):401–8.
7. McAlindon TE, LaValley MP, Harvey WF, et al. Effect of intra-articular triamcinolone vs saline on knee cartilage volume and pain in patients with knee osteoarthritis: a randomized clinical trial. *JAMA*. 2017;317(19):1967–75.
8. Dunning J, Butts R, Mourad F, et al. Dry needling: a literature review with implications for clinical practice guidelines. *Phys Ther Rev*. 2014;19(4):252–65.
9. Cagnie B, Dewitte V, Barbe T, et al. Physiologic effects of dry needling. *Curr Pain Headache R*. 2013;17(8):348.
10. Dommerholt J. Dry needling—peripheral and central considerations. *J Man Manip Ther*. 2011;19(4):223–7.
11. Gattie E, Cleland JA, Snodgrass S. The effectiveness of trigger point dry needling for musculoskeletal conditions by physical therapists: a systematic review and meta-analysis. *J Orthop Sport Phys*. 2017;47(3):133–49.
12. Brady S, McEvoy J, Dommerholt J, et al. Adverse events following trigger point dry needling: a prospective survey of chartered physiotherapists. *J Man Manip Ther*. 2014;22(3):134–40.
13. Boyce D, Wempe H, Campbell C, Fuehne S, Zylstra E, Smith G, Wingard C, Jones R. Adverse events associated with therapeutic dry needling. *Int J Sports Phys Ther*. 2020;15(1):103–13.
14. Wehling P, Evans C, Wehling J, Maixner W. Effectiveness of intra-articular therapies in osteoarthritis: a literature review. *Ther Adv Musculoskelet Dis*. 2017;9(8):183–96.
15. Gattie E, Cleland JA, Snodgrass S. A survey of American physical therapists' current practice of dry needling: practice patterns and adverse events. *Musculoskelet Sci Pract*. 2020;50:102255.
16. Naunton J, Harrison C, Britt H, Haines T, Malliaras P. General practice management of rotator cuff related shoulder pain: a reliance on ultrasound and injection guided care. *PLoS ONE*. 2020;15(1):e0227688.
17. Lin I, Wiles L, Waller R, Goucke R, Nagree Y, Gibberd M, Straker L, Maher CG, O'Sullivan PPB. What does best practice care for musculoskeletal pain look like? Eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. *Br J Sports Med*. 2020;54(2):79–86.
18. Morrissey D, Cotchett M, Said J'Barri A, Prior T, Griffiths IB, Rathlef MS, Gulle H, Vicenzino B, Barton CJ. Management of plantar heel pain: a best practice guide informed by a systematic review, expert clinical reasoning, and patient values. *Br J Sports Med*. 2021;55(19):1106–18.
19. Schneider HP, Baca JM, Carpenter BB, Dayton PD, Fleischer AE, Sachs BD. American College of Foot and Ankle Surgeons Clinical Consensus Statement: Diagnosis and Treatment of Adult acquired infra calcaneal heel pain. *J Foot Ankle Surg*. 2018;57(2):370–81. <https://doi.org/10.1053/j.jfas.2017.10.018>.
20. Martin RL, Chimenti R, Cuddeford T, Houck J, Matheson JW, McDonough CM, Paulseth S, Wukich DK, Carcia CR. Achilles pain, stiffness, and muscle power deficits: midportion Achilles tendinopathy revision 2018. *J Orthop Sports Phys Ther*. 2018;48(5):A1–38.
21. De Vos RJ, van der Vlist AC, Zwerver J, Meufels DE, Smithuis F, van Ingen R, van der Giesen F, Visser E, Balemans A, Pols M, Veen N, den Ouden M, Weir A. Dutch multidisciplinary guideline on Achilles tendinopathy. *Br J Sports Med*. 2021;29:bjsparts-2020-103867.
22. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hofmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:71
23. Shirvan R, Sadegh BM, Babak H, et al. Comparison of dry needling and steroid injection in the treatment of plantar fasciitis: a single-blind randomized clinical trial. *Int Orthop (SICOT)* 2018;42:109–6.
24. Uygur E, Aktaş B, Eceviz E. Preliminary Report on the Role of Dry Needling Versus Corticosteroid Injection, an Effective Treatment Method for Plantar Fasciitis: A Randomized Controlled Trial. *J Foot Ankle Surg* 2019;58:301–5.
25. Dr.Rahool.S, Dr.Girish.S, A prospective comparative study of local steroid Injection vs dry needling for plantar fasciitis, *IJAR*, 2022;12(1):59-63
26. Christie S, Styn G Jr, Ford G, et al. Proximal plantar intrinsic tendinopathy: anatomical and biomechanical considerations in plantar heel pain. *J Am Podiatr Med Assoc* 2019;109:412–5
27. Uygur E, Aktaş B, Yilmazoglu EG. The use of dry needling versus corticosteroid injection to treat lateral epicondylitis: a prospective, randomized, controlled study. *J Shoulder Elbow Surg* 2020;30:134–9.
28. Bhabra G, Wang A, Ebert JR, et al. Lateral elbow tendinopathy: development of a pathophysiology-based treatment algorithm. *Orthop J Sports Med* 2016;4:2325967116670635.

29. Christie S, Styn G Jr, Ford G, et al. Proximal plantar intrinsic tendinopathy: anatomical and biomechanical considerations in plantar heel pain. *J Am Podiatr Med Assoc* 2019;109:412–5.
30. Cagnie B, Dewitte V, Barbe T, et al. Physiologic effects of dry needling. *Curr Pain Headache Rep* 2013;17:348.
31. Ga H, Choi JH, Park CH, et al. Acupuncture needling versus lidocaine injection of trigger points in myofascial pain syndrome in elderly patients— a randomized trial. *Acupunct Med* 2007;25:130–6.

### **Publisher Details**

#### **SJC PUBLISHERS COMPANY LIMITED**



**Category: Non-Government & Non-profit Organisation**

**Contact: +256775434261(WhatsApp)**

**Email: [admin@sjpublisher.org](mailto:admin@sjpublisher.org), [info@sjpublisher.org](mailto:info@sjpublisher.org) or [studentsjournal2020@gmail.com](mailto:studentsjournal2020@gmail.com)**

**Website: <https://sjpublisher.org>**

**Location: Wisdom Centre Annex, P.O. BOX. 113407 Wakiso, Uganda, East Africa.**