

# A COHORT STUDY OF THE INCIDENCE OF LEFT VENTRICULAR DIASTOLIC DYSFUNCTION IN PATIENTS AFFECTED WITH SUBCLINICAL HYPOTHYROIDISM.

Nirav Kumar\*

*DM Cardio IGIMS, Patna, Bihar, India.*

---

## Abstract.

### Background:

Numerous studies have identified the presence of adverse cardiovascular effects associated with hypothyroidism. The timely detection of individuals presenting with sub-clinical hypothyroidism yields a beneficial impact on cardiovascular morbidity and mortality rates.

Objectives: The primary aim of this study is to investigate the potential correlation between subclinical hypothyroidism and left ventricular dysfunction. Additionally, we aim to explore the relationship between systolic and diastolic dysfunction in patients with subclinical hypothyroidism.

### Materials and Methods:

In this study, a total of 60 cases of subclinical hypothyroidism, as well as 30 age- and sex-matched healthy control subjects, were included. Serum levels of thyroid-stimulating hormone (TSH), thyroxine (T<sub>4</sub>), and triiodothyronine (T<sub>3</sub>) were assessed, and individuals exhibiting subclinical hypothyroidism were subsequently subjected to a two-dimensional echocardiogram (2DEcho).

### Results:

A statistically significant decrease was observed in peak early filling velocity (PE) ( $p < 0.001$ ) and early filling time velocity integral (Ei) ( $p < 0.001$ ). The statistical analysis revealed a significant reduction in the ratio of early and late peak velocities (PE/PA) ( $p < 0.001$ ), the ratio of time velocity integral of early and atrial filling (Ei/Ai) ( $p < 0.001$ ), and the ratio of the early peak to average velocity (PE/M) ( $p < 0.001$ ). The mean ejection fraction (EF) for the hypothyroid patients was  $54.8 \pm 5.54$ , which was slightly lower compared to the control subjects who had a mean EF of  $55.6 \pm 3.45$ . The difference between the two groups was not statistically significant, as indicated by a T-value of 0.48.

### Conclusion:

Our findings indicate that sub-clinical hypothyroidism is associated with notable diastolic dysfunction, while systolic function remains largely unaffected.

### Recommendation:

Patients with subclinical hypothyroidism should be regularly monitored for their diastolic function. Any abnormality if detected should be promptly managed to reduce cardiovascular morbidity

**Keywords:** Left ventricular diastolic dysfunction, Subclinical hypothyroidism, Cardiac dysfunction, Diastolic, Echocardiographic evaluation, Submitted: 2023-08-30 Accepted: 2023-09-10

---

## 1. INTRODUCTION.

The cardiac manifestations observed in individuals with subclinical hypothyroidism primarily consist of systolic dysfunction [1]. The identification of adverse cardiovascular effects associated with hypothyroidism has been well-documented in numerous studies. The majority of the investigations have primarily concentrated on the assessment of systolic and diastolic dysfunction in individuals with hypothyroidism [2-6]. Subclinical hypothyroidism (SCH) is characterized by an elevation in serum thyroid stimulating hormone (TSH) levels that surpass the upper threshold of normal while maintaining normal levels of serum-free thyroxine. The prevailing etiology of this condition is commonly attributed to autoimmune thyroid disease [7]. Subclinical hypothyroidism (SCH) is a frequently encountered medical condition, exhibiting a prevalence ranging from 3% to 8% among individuals without a documented history of thyroid pathology. The prevalence of the condition increases proportionally with advancing age and exhibits a higher incidence among the female population. The presence of antithyroid antibodies has been observed in approximately 80% of patients diagnosed with subclinical hypothyroidism (SCH). In the context of subclinical hypothyroidism (SCH), it is observed that approximately 80% of patients exhibit a serum thyroid-stimulating hormone (TSH) level below  $10 \mu\text{IU/L}$  [8]. Subclinical hypothyroidism (SCH) has the potential to advance into overt hypothyroidism, thereby giving rise to systemic manifestations of hypothyroidism. These manifestations include cardiac dysfunction, adverse effects on fetal development, neuromuscular impairments, psychiatric disturbances, cognitive dysfunction, elevation in serum total cholesterol levels, as well as increased levels of low-density lipoprotein cholesterol (LDL-C) [9]. The impact of SCH on the cardiovascular system has been evaluated through the examination of diastolic function, characterized by decelerated myocardial relaxation and compromised

ventricular filling. The presence of this cardiac abnormality may potentially hurt the prognosis due to its association with isolated left ventricular diastolic dysfunction (LVDD), which is correlated with heightened morbidity and mortality rates in the overall population [10]. Diastolic dysfunction possesses crucial therapeutic implications and can aid physicians in formulating prompt intervention strategies [11]. The yearly probability of advancing to overt hypothyroidism is estimated to be around 5% in cases where subclinical hypothyroidism (SCH) is present alongside elevated levels of auto-antibodies in the bloodstream. Subacute combined hyperhomocysteinemia (SCH) manifests with a clinical presentation that lacks specificity and is commonly confirmed through laboratory investigations [12]. The assessment of diastolic dysfunction can be accomplished through the utilization of pulsed-wave Doppler (PWD) for the measurement of mitral inflow velocities and tissue Doppler echocardiography (TDE) for the measurement of mitral annular velocities [13]. The timely detection of individuals presenting with sub-clinical hypothyroidism holds the potential to initiate prompt intervention, consequently yielding a beneficial impact on cardiovascular morbidity and mortality rates. The present study has been designed to investigate the impact of sub-clinical hypothyroidism on left ventricular diastolic functions.

### 1.1. Objectives.

- *The objective of this study is to investigate the potential correlation between subclinical hypothyroidism and left ventricular dysfunction.*
- *The objective of this study is to investigate the correlation between systolic and diastolic function in the selected patient population.*

## 2. MATERIALS AND METHODS.

The cohort study was carried out within the Non-invasive cardiology laboratory situated at IGIMS, Patna. The inclusion criteria for the

\*Corresponding author.

Email address: [drniravsinha22@gmail.com](mailto:drniravsinha22@gmail.com) (Nirav Kumar)

study was subclinical hypothyroidism so it encompassed a total of 60 cases of subclinical hypothyroidism, accompanied by 30 age- and sex-matched healthy control subjects. The study exclusively encompassed cases in which the diagnosis of subclinical hypothyroidism was established using standard criteria. Exclusion criteria encompassed various conditions that have been associated with left ventricular dysfunction, including but not limited to advanced age (greater than 70 years), elevated heart rate (above 100 beats per minute), conditions leading to left ventricular hypertrophy, conditions resulting in myocardial stiffness such as hemochromatosis, amyloidosis, alcoholism, diabetes mellitus, sarcoidosis, and other infiltrative diseases. Additionally, individuals with overt hypothyroidism and the presence of any chronic medical or surgical condition that may independently detrimentally impact left ventricular function were also excluded from the study. Serum levels of thyroid-stimulating hormone (TSH), thyroxine (T<sub>4</sub>), and triiodothyronine (T<sub>3</sub>) were assessed, and individuals exhibiting subclinical hypothyroidism were subsequently subjected to a two-dimensional echocardiogram (2DEcho). Informed consent was obtained from all patients before they participated in the study.

### 3. RESULTS.

The patients who had subclinical hypothyroidism were included in the study and those who met the exclusion criteria specified were eliminated. In all, there were 60 eligible patients for the study. The average age of patients was  $46.9 \pm 6.92$  years, whereas, for the control group, it was  $47.4 \pm 5.96$  years. In a cohort of 60 cases, the proportion of female patients was 76.66%, while the proportion of male patients was 23.33%. Out of the total sample size of 54 cases, representing 90% of the population, the majority exhibited no clinical manifestations indicative of hypothyroidism. However, a small subset of 2 cases (3.3%) presented with a dull expression, while another 2 cases displayed sparse hair. Additionally, 6 cases (10%) were found to be associated with prolonged relaxation of the Achilles tendon re-

flex. No electrocardiographic abnormalities were observed in any of the patients. In the present study, a total of 52 individuals (86.66%) exhibited no evidence of dyslipidemia, while 8 cases (13.3%) demonstrated an association with elevated levels of total cholesterol. Additionally, 4 cases (6.6%) were found to be associated with increased levels of triglycerides, as depicted in Table 1.

The Doppler evaluation reveals a substantial decrease in the peak early filling velocity (PE) ( $p < 0.001$ ) and early filling time velocity integral (Ei) ( $p < 0.001$ ), indicating a significant impairment in cardiac diastolic function. The ratio of early and late peak velocities (PE/PA) demonstrated a statistically significant decrease ( $p < 0.001$ ). Similarly, the ratio of time velocity integral of the early and atrial filling (Ei/Ai) exhibited a significant reduction ( $p < 0.001$ ). Additionally, the ratio of the early peak to average velocity (PE/M) was found to be significantly decreased ( $p < 0.001$ ). The mean ejection fraction (EF) of the hypothyroid patients was  $54.8 \pm 5.54$ , which was compared to the EF of the control subjects, which was  $55.6 \pm 3.45$ . The difference between the two groups was not statistically significant, with a T value of 0.48. However, there was a significant presence of diastolic dysfunction in the hypothyroid patients, as indicated by the mean Ei/Ai ratio of  $1.34 \pm 0.52$ . In comparison, the control group subjects had a mean Ei/Ai ratio of  $2.12 \pm 0.27$ . The difference between the two groups in terms of the Ei/Ai ratio was highly statistically significant, with a T value of 5.22 (Table 2).

### 4. DISCUSSION.

Conflicting evidence exists concerning the phenomenon of subclinical hypothyroidism. According to a study, it has been determined that subclinical hypothyroidism, when left undetected, does not pose a risk factor for cardiovascular health [14]. The study on cardiovascular health revealed no discernible elevation in cardiovascular risk among individuals with thyroid-stimulating hormone (TSH) levels below 10.0 milliunits per liter (mU/l) [15]. In a separate investigation,

Table 1: Study population characteristics.

	Controls	Patients
<b>Mean age* (<math>\pm</math> SD) Years</b>	47.4 $\pm$ 5.96	46.9 $\pm$ 6.92
<b>Sex</b>		
Male	10 (33.33)	14 (23.33)
Female	20 (66.66)	46 (76.66)
<b>Signs/Symptoms</b>		
Constipation	-	6 (10)
Lethargy	-	6 (10)
Menstrual disorder	-	6 (10)
Increase in weight	-	6 (10)
Prolonged relaxation of deep tolerance reflex	-	6 (10)
Decreased appetite	-	4 (6.66)
Sparse hair	-	2 (3.33)
Dull expressionless face	-	2 (3.33)
Cold intolerance	-	2 (3.33)
<b>Lipid profile</b>		
HDL cholesterol (<66 mg%)	-	0 (0.0)
LDL cholesterol (>155 mg%)	-	0 (0.0)
Total cholesterol (>250 mg%)	-	8 (13.3)
Triglyceride (>180 mg%)	-	4 (6.6)
Within normal limit	30 (100)	52 (86.66)

the study postulated that patients exhibiting a thyroid-stimulating hormone (TSH) level exceeding 7 mIU/L may potentially face an elevated susceptibility to heart failure. However, no other notable cardiovascular incidents were observed in this cohort [16]. Based on the data that has been reviewed, it is recommended to refrain from implementing routine treatment for subclinical hypothyroid disease in cases where the thyroid-stimulating hormone (TSH) level falls within the range of 4.5–10.0 mIU/L [12]. In contrast, Biondi et al. discovered an association between subclinical hypothyroidism and left ventricular systolic and diastolic dysfunction, as well as an increased risk for atherosclerosis and myocardial infarction. This association is based on the observation that the heart exhibits a response to minimal, yet persistent alterations in thyroid hormone levels, which are characteristic of subclinical hypothyroidism [17]. There is also compelling evidence

about the heightened arterial stiffness observed in this pathological condition [18]. Several studies have provided evidence of diastolic dysfunction in individuals with subclinical hypothyroidism. These findings have led researchers to suggest that treatment with levothyroxine may potentially alleviate this diastolic dysfunction over a certain duration of time [19–21].

A research study has revealed the presence of notable diastolic dysfunction in the left ventricle among patients diagnosed with subclinical hypothyroidism, while the systolic function remains unaltered. A notable augmentation in mitral peak A velocity, diminished E/A ratio, and elongation of isovolumic relaxation time (IVRT) were observed in the Doppler echocardiography findings. The mitral peak E velocity remains unchanged. Subclinical hypothyroidism exhibits a notably elevated Tei index in comparison to individuals with normal thyroid function. The present obser-

Table 2: Comparative analysis was conducted to assess the left ventricular diastolic and systolic function in two distinct groups: control subjects and sub-clinical hypothyroid patients.

	<b>Indices of LV function</b>	<b>Patient</b>	<b>Control</b>	<b>t</b>	<b>P</b>
<b>LV Systolic function</b>		(mean ± SD)	(mean ± SD)		
1	EF	54.8 ± 5.54	55.6 ± 3.45	-0.48	
<b>LV Diastolic function</b>	<b>Indices of LV function</b>	<b>Patient</b>	<b>Control</b>	<b>t</b>	<b>P</b>
		(mean ± SD)	(mean ± SD)		
1	PE	0.51 ± 0.11	0.65 ± 0.07	-4.55	<0.001
2	PA	0.53 ± 0.11	0.52 ± 0.08	-0.05	
3	M	0.33 ± 0.09	0.30 ± 0.06	0.32	
4	Ei	4.21 ± 0.79	7.53 ± 1.61	-6.19	<0.001
5	Ai	3.50 ± 1.48	3.60 ± 0.77	-0.29	
6	Ti	9.78 ± 2.36	13.16 ± 2.91	-3.15	<0.01
7	%AC	35.54 ± 10.26	27.58 ± 2.84	3.23	<0.01
8	1/3FF	38.23 ± 6.67	47.02 ± 4.18	-4.41	<0.001
9	P%T	0.09 ± 0.04	0.12 ± 0.02	-2.78	<0.01
10	Time E	0.09 ± 0.02	0.11 ± 0.03	-3.04	<0.01
11	PE/PA	0.98 ± 0.23	1.28 ± 0.15	-3.87	<0.001
12	PE/M	1.64 ± 0.40	2.28 ± 0.36	-4.29	<0.001
	Ei/Ai	1.34 ± 0.52	2.12 ± 0.27	-5.22	<0.001

vations bear resemblance to the aforementioned findings reported by Biondi et al., albeit without the inclusion of Tei index measurements [22]. The present study provides additional evidence that aligns with the findings of a study., indicating a notable decrease in the E/A ratio and elongation of the isovolumic relaxation time (IVRT) in both the left and right ventricles. However, it is important to note that our study did not encompass an assessment of right ventricular dysfunction [23]. The incidence of cardiovascular disease (CVD) has been documented to be lower in the female population compared to males before the onset of menopause. This disparity is believed to be associated with a diminished accumulation of abdominal adipose tissue in women [24]. The mean body mass index (BMI) observed in our

study was slightly elevated, approaching the upper threshold of the normal range, consistent with findings reported in previous studies [10, 25, 26]. A notable correlation has been observed between body mass index (BMI) and subclinical hypothyroidism (SCH) [27]. SCH, also known as subclinical hypothyroidism, is a thoroughly characterized medical condition exhibiting a comprehensive array of clinical manifestations and symptoms [28]. The presence of diastolic dysfunction has been associated with a higher prevalence of diabetes, hypertension, and dyslipidemia in patients [29]. In the presence of abnormal myocardial relaxation, there is an observed increase in the ratio of mitral annular motion during atrial systole to total diastolic annular motion [30]. The precise characterization of the impact of modified diastolic function

in individuals with compromised systolic function remains to be definitively established, albeit its potential significance cannot be understated. Primary diastolic dysfunction, in the absence of systolic dysfunction, represents a significant and progressively acknowledged medical condition.

## 5. CONCLUSION.

In our study, it was observed that sub-clinical hypothyroidism exhibited notable diastolic dysfunction while not significantly affecting systolic function, as evidenced by the measurement of Ejection Fraction.

## 6. RECOMMENDATION.

Patients with sub clinical hypothyroidism should be regularly monitored for their diastolic function. Any abnormality if detected should be promptly managed to reduce cardiovascular morbidity.

## 7. LIMITATION.

This was a hospital-based study carried out on a small population, more inclusive studies on a large population are required to conclude the effects of hypothyroidism on the cardiovascular system.

## 8. LIST OF ABBREVIATIONS.

TSH- Thyroid-stimulating hormone  
2D-echo-two-dimensional echocardiogram  
PWD- pulsed-wave Doppler  
SCH- Subclinical hypothyroidism  
LVDD-left ventricular diastolic dysfunction  
TDE- tissue Doppler echocardiography  
IVRT- isovolumic relaxation time  
CVD-cardiovascular disease  
BMI- body mass index

## 9. ACKNOWLEDGEMENT.

We are thankful to the patients and their caring parents without them the study could not have been done. We are thankful to the supporting staff of our hospital who were involved in the patient care of the study group.

## 10. Conflict of interest.

The study was not funded.

## 11. Funding.

The study was not funded.

## 12. Publisher details:

**Publisher: Student's Journal of Health Research (SJHR)**  
**(ISSN 2709-9997) Online**  
**Category: Non-Governmental & Non-profit Organization**  
**Email: [studentsjournal2020@gmail.com](mailto:studentsjournal2020@gmail.com)**  
**WhatsApp: +256775434261**  
**Location: Wisdom Centre, P.O.BOX. 148, Uganda, East Africa.**



## 13. REFERENCES.

1. Bionadi B, Klein I, et al. Hypothyroidism as a risk factor for cardiovascular disease. *Endocrine*. 2004;24(1):1–13.
2. Shenoy MM, Goldman JM. *American Journal of Medical Sciences*. 1987;294:1–9.
3. o'Malley S, Peterson A, Mc Cullough FD, Rosenthal, et al. Reversible abnormalities of myocardial relaxation in hypothyroidism. *J Clin Endocrinol Metab*. 1985;61:269–272.

4. Farfar JC, Muir AL, Toft AD. Left ventricular function in hypothyroidism response to exercise and beta-adrenoceptor blockade. *Br Heart J*. 1982;48:278–284.
5. William F Crowley, Chester Ridgway, et al. Non invasive evaluation of cardiac function in hypothyroidism. *New England Journal*. 1977;296:1–6.
6. Bough EW, Crowley WF, et al. Myocardial function in hypothyroidism. *Arc Intern Med*. 1978;138:1476–1480.
7. Cooper DS. Subclinical hypothyroidism. *N Engl J Med* 2001;345:260–265.
8. Hollowell JG, Staehling NW, Flanders WD, et al. Serum TSH, T4, and thyroid antibodies in the United States population: National health and nutrition examination survey. *J Clin Endocrinol Metab* 2002;87:489–499.
9. Chu JW, Crapo LM. The treatment of subclinical hypothyroidism is seldom necessary. *J Clin Endocrinol Metab* 2001;86:4591–4599.
10. Akcakoyun M, Kaya H, Kargin R, et al. Abnormal left ventricular longitudinal functional reserve assessed by exercise pulsed wave tissue Doppler imaging in patients with subclinical hypothyroidism. *J Clin Endocrinol Metab* 2009;94:2979–2983.
11. Meena CL, Meena RD, Nawal R, et al. Assessment of left ventricular diastolic dysfunction in subclinical hypothyroidism. *Acta Inform Med* 2012;20:218–220.
12. Hak E, Pols HAP, Visser TJ. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and MI in elderly women. *Ann Intern Med* 2000;132:270–278.
13. Sohn DW, Chai IH, Lee DJ, et al. Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic dysfunction. *J Am Coll Cardiol* 1997;30:474–480.
14. Cappola AR, Fried LP, Arnold AM, Danese MD, Kuller LH, Burke GL, et al. Thyroid status, cardiovascular risk, and mortality in older adults. *JAMA* 2006;295:1033–41.
15. Rodondi N, Bauer DC, Cappola AR, Cornuz J, Robbins J, Fried LP, et al. Subclinical thyroid dysfunction, cardiac function, and the risk of heart failure. The cardiovascular health study. *J Am Coll Cardiol* 2008;52:1152–9.
16. Rodondi N, Newman AB, Vittinghoff E, de Rekeneire N, Satterfield S, Harris TB, et al. Subclinical hypothyroidism and the risk of heart failure, other cardiovascular events, and death. *Arch Intern Med* 2005;165:2460–6.
17. Biondi B, Palmieri EA, Lombardi G, Fazio S. Effects of subclinical thyroid dysfunction on the heart. *Ann Intern Med* 2002;137:904–14.
18. Owen PJ, Rajiv C, Vinereanu D, Mathew T, Fraser AG, Lazarus JH. Subclinical hypothyroidism, arterial stiffness, and myocardial reserve. *J Clin Endocrinol Metab* 2006;91:2126–32.
19. Vitale G, Galderisi M, Lupoli GA, Celentano A, Pietropaolo I, Parenti N, et al. Left ventricular myocardial impairment in subclinical hypothyroidism assessed by a new ultrasound tool: Pulsed tissue Doppler. *J Clin Endocrinol Metab* 2002;87:4350–5.
20. Franzoni F, Galetta F, Fallahi P, Tocchini L, Merico G, Braccini L, et al. Effect of L-thyroxine treatment on left ventricular function in subclinical hypothyroidism. *Biomed Pharmacother* 2006;60:431–6.
21. Erkan G, Erkan AF, Cemri M, Karaahmetoglu S, Cesur M, Cengel A. The evaluation of diastolic dysfunction with tissue Doppler echocardiography in women with subclinical hypothyroidism and the effect of L-thyroxine treatment on diastolic dysfunction: A pilot study. *J Thyroid Res* 2011;2011:654304.
22. Biondi B, Fazio S, Palmieri EA, Carella C, Panza N, Cittadini A, et al. Left ventricular diastolic dysfunction in patients with subclinical hypothyroidism. *J Clin Endocrinol Metab* 1999;84:2064–7.
23. Kosar F, Sahin I, Turan N, Topal E, Aksoy Y, Taskapan C. Evaluation of right and left ventricular function using pulsed-wave tissue Doppler echocardiography in patients with subclinical hypothyroidism. *J Endocrinol Invest* 2005;28:704–10.

24. Van Pelt RE, Jankowski CM, Gozansky WS, et al. Sex differences in the association of thigh fat and metabolic risk in older adults. *Obesity (Silver Spring)* 2011;19:422–428.
25. Tadic M, Ilic S, Kostic N, et al. Sub-clinical hypothyroidism and left ventricular mechanics: A three-dimensional speckle tracking study. *J Clin Endocrinol Metab* 2014;99:307–314.
26. Pesic M, Antic S, Kocic R, et al. Cardiovascular risk factors in patients with sub-clinical hypothyroidism. *Vojnosanit Pregl* 2007;64:749–752.
27. Meher LK, Raveendranathan SK, Kota1 SK, et al. Prevalence of hypothyroidism in patients with metabolic syndrome. *Thyroid Res Pract* 2013;10:60–64.
28. McDermott MT, Ridgway EC. Subclinical hypothyroidism is mild thyroid failure and should be treated. *J Clin Endocrinol Metab* 2001;86: 4585–4590.
29. Paul R, Sinha PK, Mookherjee S, et al. Prevalence of left ventricular diastolic dysfunction in patients with normal systolic function and its relation with other risk factors from eastern India. *Ind Med Gaz* 2015; 2:54–61.
30. Vitale G, Galderisi M, Lupoli GA, et al. Left ventricular myocardial impairment in sub-clinical hypothyroidism assessed by a new ultrasound tool: pulse tissue Doppler. *J Clin Endocrinol Metab* 2002;87:4350–4355.

### **Author biography**

**Nirav Kumar** DM Cardio, Associate Professor, IGIMS, Patna, Bihar, India